



Attack-All[®]

LIVESTOCK & PREMISE FLY SPRAY

- **READY TO USE**
- **KILLS & REPELS: HOUSE FLIES, STABLE FLIES, FACE FLIES, HORN FLIES, GNATS AND MOSQUITOES**
- **USE ON CATTLE (BEEF & DAIRY) OR SWINE**
- **APPLY TO: MILKING PARLORS, CALF HUTCHES, BARNs, POULTRY HOUSES, SWINE QUARTERS, PENS & SHELTERS**

SPECIMEN LABEL

ACTIVE INGREDIENT:

Pyrethrins	0.05%
Piperonyl Butoxide*	0.50%
Permethrin**	0.10%
OTHER INGREDIENTS	99.35%
TOTAL	100.00%

* (butylcarbityl)(6-propylpiperonyl) ether and related compounds
 ** cis/trans ratio: min. 35% (+/-) cis and max 65% (+/-) trans

EPA Reg. No. 2724-810

EPA Est. No. 2724-TX-1

KEEP OUT OF REACH OF CHILDREN
CAUTION

See additional Precautions

PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Not for use on humans. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Hazards to Domestic Animals: For external use only. Do not use on animals under three months of age or on sick, old, debilitated, or medicated animals.

FIRST AID

IF ON SKIN • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 – 20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-347-8272 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Drift and runoff may be hazardous to fish in water adjacent to treated areas. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. See Directions for Use for additional precautions and requirements. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON BEEF AND DAIRY CATTLE OR SWINE.

Starbar[®] Attack-All[®] Livestock & Premise Fly Spray (Attack-All[®]) can be used on cattle, swine and in animal quarters as a fast, effective product for control of flying insects.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not remain in treated area. Exit area immediately and remain outside the treated area until mists have dispersed.

Do not allow adults, children, or pets to enter until sprays have dried or dispersed and the treated area has been thoroughly ventilated.

Do not apply during rain.

Do not water the treated area to the point of runoff.

Remove or cover exposed food and drinking water before application. Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces or wash them before use.

All outdoor applications must be limited to spot or crack-and-crevice treatments only except for the following permitted uses:

- (1) Treatment to soil or vegetation around structures;
- (2) Applications to building foundations, up to a maximum height of three feet.

As a space spray, the maximum use rate is 2 fluid ounces per 1,000 cubic feet (10 ft x 10 ft x 10 ft high) of space. For space sprays, do not apply more than one time per day. On animals, the maximum use rate is 3 fluid ounces per animal at a maximum of one application per ten days (cattle) or per 14 days (swine).

ON LIVESTOCK (Indoors and Outdoors): To protect beef cattle, dairy cattle, and swine from house flies, horn flies, face flies, gnats, and mosquitoes, apply a light mist sufficient to wet the surface of the hair. For cattle, do not reapply for ten days. For swine, do not reapply for 14 days. For swine, do not apply five days or less prior to slaughter.

DIRECTIONS FOR TRIGGER SPRAY USE: Remove excess dirt and dust. Apply light spray mist to coat while brushing lightly against lay of the hair. Avoid spraying in eyes and mucous membranes. Apply with sponge or cloth to those areas. Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.

TO CONTROL stable flies, horse flies, and deer flies, apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment at ten day intervals for cattle or 14 day intervals for swine to give continued protection.

To protect from face flies, spray on a cloth and rub on the face, being certain to avoid contact with eyes and mucous membranes.

TO CONTROL blood-sucking lice, apply to infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every two to three weeks if required.

TO KILL AND REPEL FLYING INSECTS IN ANIMAL QUARTERS: Use this product in cattle (beef and dairy) barns, milk houses, poultry houses, swine quarters, outbuildings, pens and shelters to kill and repel stable flies, horse flies, face flies, deer flies, house flies, horn flies, mosquitoes, gnats, ticks (including deer ticks), fleas, chiggers, and lice. Apply undiluted as either a surface spray or space spray (fog). Remove animals prior to treatment. When used in dairy barns or facilities, close milk bulk tank lids to prevent contamination.

Surface Spraying: Use a good sprayer adjusted to deliver a coarse, wet spray. Apply as a coarse, wet spray using 1 pint per 1,000 square feet. Direct the spray wherever flies congregate or alight, until surfaces are moistened. Treat around doorways, feed storage rooms, alleyways, and windowsills. Spray around outside of door facings and screens to render area unattractive to insects. This helps to repel insects and prevent entrance into building.

Space Spraying: Before application, close doors, windows, and other openings and shut off ventilating systems to reduce air movement as much as possible. Apply with a good machine adjusted to deliver a fine mist or fog. Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 2 fluid ounces per 1,000 cubic feet (10 ft x 10 ft x 10 ft). Apply by directing

spray toward ceiling and corners until the space is filled with mist. For best results, keep doors and windows closed, when possible, during spraying and after application. Applicator must not remain in treated area. Leave room closed for ten minutes after application, then open and ventilate the area thoroughly prior to reoccupation by humans or animals.

Do not contaminate water, feed or foodstuffs, milk or milking utensils. Repeat treatment after ten days, or at intervals needed to give adequate protection.

TO CONTROL poultry lice, spray roosts, walls, and nests or cages thoroughly. Follow by spraying over the birds with a fine mist.

FOR CONTROL of bedbugs and mites in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

OUTDOOR USE: Use this product to control cockroaches, waterbugs, palmetto bugs, scorpions, millipedes, centipedes, sowbugs, pillbugs, ants, silverfish, firebrats, spiders, and crickets. To help prevent infestation of building, treat the building foundation to a height of two to three feet, where pests are active and may find entrance. Also apply as a residual spray to outside surfaces of buildings, and other areas where these pests congregate or have been seen.

Avoid spraying when wind conditions might create a mist that can blow back to applicator. Do not apply this product in or around bodies of water. Do not spray directly on plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store in a cool, dry place. Do not store near heat or open flame. **CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY AND CONDITIONS OF SALE

Seller warrants that this product conforms to the chemical specifications expressed on the label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, seller's liability for default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

For information or in case of an emergency, call 1-800-347-8272.

www.starbarproducts.com

Manufactured for:

Wellmark International
1501 East Woodfield Road 200W
Schaumburg, Illinois 60173

